

# Physical and nutritional properties of hawthorn fruit (*Crataeguspontica* L.)

JavadErfani Moghadam<sup>1\*</sup>, Kamran Kheiralipour<sup>2</sup>

(1. Department of Horticulture, Ilam University, Ilam, I.R. Iran 69315-516

2. Department of Mechanical Engineering of Agricultural Machinery, Ilam University, Ilam, I.R. Iran 69315-516)

**Abstract:** Hawthorn, from Rosaceae family, is one of the important forest fruits of which there are different species in Iran. In this study, some characteristics of hawthorn fruit (*Crataeguspontica*), including physical properties (fruit length, width, thickness, geometric, arithmetic and equivalent mean diameter, surface area, sphericity, aspect ratio, thousand fruit mass and true density) and nutritional properties (total dry matter, total soluble solid, titratable acidity and moisture content) were considered. Results showed that average of fruit length, width and thickness were 1.53 mm, 1.95 mm and 1.78 mm, respectively. The geometric mean diameter was same to arithmetic mean diameter (1.75mm), while equivalent mean diameter was higher than both (1.76 mm). Some physical properties such as sphericity (1.13%), surface area (1.69mm) and aspect ratio (1.26) were determined. Average of 100 fruit weight in this species was 306.54 g and it is estimated 3.06 g for one fruit. Total soluble solid percent (TSS) and titratable acidity percent (TA) of fruit hawthorn were estimated 18.7% and 1.71%, respectively.

**Keywords:** hawthorn fruit, physical characteristics, nutritional properties

**Citation:** Erfani, J. M., and K. Kheiralipour. 2015. Physical and nutritional properties of hawthorn fruit (*Crataeguspontica* L.). *AgricEngInt: CIGR Journal*, 17(1): 232-237.

## 1 Introduction

Physical properties of agricultural products are useful information respect formechanizing harvest and postharvest operations. One of these properties is physical characteristics including information about dimension and mass. Nutritional properties are important to know how a fruit or vegetable can be considered as potential in food industry.

Numerous researches were conducted to collect physical information of different fruits and vegetables such as: sunflower seeds (Perez et al., 2007), cowpea seed (Kabas et al., 2007), cucurbit seeds (Milani et al., 2007),

watermelon seed (Koochehi et al., 2007), rice (Ghasemi Varnamkhasti et al., 2008), apple (Kheiralipour, 2008; Ragni et al., 2011), pomegranate (Riyahi et al., 2011), almond nut and kernel (Mirzabe et al., 2013) and wild pistachio (Fadavi et al., 2013).

In this research, hawthorn fruit of Ilam, Iran was considered. Hawthorn includes more than 250 different species (Blumenthal et al., 2000). The varieties are from small shrubs to trees (Baharun et al., 2003). In general, the forms of the varieties are not as large trees or canopy dominants. Some of them are decidedly shrubby, whereas others can grow largely to heights of 12 m (Kumar et al., 2012). Hawthorn is distributed well in western Asia, Europe and North America. It is widely used horticultural as an ornamental. The colors of ripped hawthorn fruits are yellow, red, black and the combined colors (Donmez, 2004).

Received date: 2014-09-21 Accepted date: 2014-12-22

\*Corresponding author: JavadErfaniMoghadam, Department of Horticulture, Ilam University, Ilam, I.R. Iran 69315-516. Tel & Fax: +98-841-2227015. E-mail address: j.erfani@ilam.ac.ir

Another use of hawthorn is pharmaceutical utility in the western world (Bahorun et al., 2003; Ercisli, 2004). It is considered as one of the oldest traditional medicinal plants in some countries (Blumenthal et al., 2000) due to numerous chemical constituents (Ozturk and Tuncel, 2011). The medicinal aspects of hawthorn have a long tradition according to records written in ancient Roman period (Veveris et al., 2004). Some valuable antioxidants are predicted to find as some *Crataegus* components. The fruit and also the flower components of hawthorn are responsible for free radical scavenging activities such as epicatechin and chlorogenic acid. They can be well among the best anti-lipoperoxidants (Güven et al., 2006).

Iran is recognized as an important source of genetic diversity of different plant species including fruit crops. Also, the country is a rich source of genetic variability for *Crataegus* species germplasm, which have a high morphological diversity, particularly in leaf, flower, and fruit characteristics. The hawthorn fruits of Ilam province are yellow to yellowish red and black, and those of Kurdistan province are red. In the present study, the authors describe some fruit attributes of *Crataegus pontica* grown in the Ilam forests including physical and nutritional properties. These properties can be useful in designing the machines for harvest, post-harvest processing operations and further breeding efforts.

## 2 Materials and methods

Hawthorn fruit (*Crataegus pontica*) were randomly harvested from forests of Ilam province in late Oct. 2012 (Figure 1). Experiments were conducted in randomized block design with three repetitions and a total of 100 fruits for each repeat. The research was carried out in Ilam University, Ilam, Iran.



Figure 1 Hawthorn, fruit and kernel

## 3 Physical characteristics

Considered physical characteristics of hawthorn fruit were, length, width, thickness, geometric diameter, equivalent diameter, arithmetic diameter, sphericity, surface area, aspect ratio, volume, bulk density, true density and porosity.

To determine the average size of the fruits, three linear dimensions, namely length,  $L$ , width,  $W$ , and thickness,  $T$ , were measured by using a digital caliper with accuracy of 0.01 mm, and fruit mass was determined with an electronic balance of 0.1 g accuracy. The diameters were calculated by considering Equation 1, Equation 2 and Equation 3, respectively (Mohsenin, 1986):

$$D_g = (LWT)^{\frac{1}{3}} \quad (1)$$

$$D_p = \left[ L \frac{(W+T)^2}{4} \right]^{\frac{1}{3}} \quad (2)$$

$$D_a = \frac{(L+W+T)}{3} \quad (3)$$

where:

$L$ : Fruit length (mm)

$W$ : Fruit width (mm)

$T$ : Fruit thickness (mm)

$D_g$ : Geometric diameter (mm)

$D_p$ : Equivalent diameter (mm)

$D_a$ : Arithmetic diameter (mm).

The sphericity,  $Sp(\%)$ , is the ratio of surface area of a sphere having the same volume as that of fruit to the surface area of the fruit, was determined using the following Equation 4 (Mohsenin, 1986):

$$D_p = \frac{(LWT)^{\frac{1}{3}}}{L} \quad (4)$$

The surface area,  $S$  ( $\text{mm}^2$ ), of the fruit was calculated by using the following Equation 5 (Mohsenin, 1986):

$$S = \frac{\pi}{4} D_g^2 \quad (5)$$

The aspect ratio,  $R_a$  was calculated by Equation 6 (Omobuwajo et al. (1999):

$$R_a = \frac{W}{L} \quad (6)$$

To determine volume, bulk density, true density and porosity, a few fruits with specific masses were dropped on container with volume of  $200 \text{ cm}^3$ . The container was full of water to fill the porous spaces. The fruit volume was calculated by Equation 7:

$$V_f = V_c - V_w \quad (7)$$

Where:

$V_f$ : Fruit volume ( $\text{cm}^3$ )

$V_c$ : volume of container ( $\text{cm}^3$ )

$V_w$ : water volume ( $\text{cm}^3$ ).

Bulk density and true density were calculated using Equation 8 and Equation 9, respectively.

$$\rho_b = \frac{M_f}{V_c} \quad (8)$$

$$\rho_t = \frac{M_f}{V_f} \quad (9)$$

Where:

$\rho_b$ : Bulk density ( $\text{g}/\text{cm}^3$ )

$\rho_t$ : True density ( $\text{g}/\text{cm}^3$ )

$M_f$ : Fruit mass (g).

The fruit porosity,  $\varepsilon$  (%), was determined by following Equation 10 (Mohsenin, 1986):

$$\varepsilon = \frac{V_w}{V_f} \quad (10)$$

## 4 Nutritional properties

Nutritional properties of the fruit were included, titratable acidity (TA), total soluble solids (TSS), TSS/TA ratio, dry matter and moisture content. Three samples of fruit texture (10 g) were kept in an oven in  $70 \text{ }^\circ\text{C}$  for 48 h. Then, the moisture content and dry matter were calculated by Equation 11 and Equation 12, respectively:

$$MC = \frac{M_w - M_d}{M_w} \times 100 \quad (11)$$

$$DM = \frac{M_d}{M_w} \times 100 \quad (12)$$

Where:

$M_w$ : Initial mass of fruit (g)

$M_d$ : Mass of dried fruit (g)

MC: Moisture content of fruit, wet base (%)

DM: fruit Dry matter (%).

Titratable acidity (TA) was determined by potentiometric titration with 0.1 N NaOH up to pH 8.2, using 5 ml of juice diluted to 50 ml with distilled  $\text{H}_2\text{O}$ . The results were expressed as gram of malic acid per 100 g fresh weight. Total soluble solids (TSS, °Brix) were determined using a hand-held refractometer, also on ten fruit. TSS/TA ratio was specified as following Equation 13:

$$T = \frac{TSS}{TA} \quad (13)$$

Where:

T: TSS/TA ratio

TSS: Total soluble solids (°Brix)

TA: Titratable acidity (%).

## 5 Results and discussion

The values of all measured and calculated physical characteristics of Ilam hawthorn fruit were shown in Table 1.

**Table 1 Physical characteristics of hawthorn fruit**

Parameter	Maximum	Mean	Minimum	Standard Deviation
Length, mm	18.20	15.47	13.00	1.23
Width, mm	22.5	19.50	15.53	1.64
Thickness, mm	20.40	17.78	14.30	1.58
Geometric mean diameter, mm	19.84	17.51	14.62	1.37
Equivalent diameter, mm	19.86	17.53	14.63	1.37
Arithmetic diameter, mm	19.93	17.60	14.63	1.38
Sphericity, %	1.24	1.13	1.04	0.04
Surface area, mm <sup>2</sup>	1235.51	968.41	671.20	150.10
Aspect ratio	1.39	1.26	1.10	0.07
Thousand fruit weight, g	3104.86	3065.47	2988.55	66.62
True density, kg/m <sup>3</sup>	0.93	0.89	0.83	0.05
Bulk density, kg/m <sup>3</sup>	0.48	0.47	0.46	0.01
Porosity, %	48.50	47.50	46.50	0.01

The mean of fruit length (L), width (W), and thickness (T) were 15.47, 19.50, and 17.88 mm, respectively. Geometric mean diameter and equivalent diameter of the fruit were close (17.51 and 17.53, respectively) whereas the arithmetic diameter had slightly higher magnetite (17.60). The critical use of fruit dimensional properties is to determine the aperture size of machines, particularly in separation of materials (Mohsenin, 1978). In this research, the dimensions of the fruits are relatively uniform hence it will improve the process optimization in design and development harvesting machine for the fruit as well as in separation and cleaning.

Other dimensional properties were 1.13%, 968.41 mm<sup>2</sup> and 1.26 as sphericity, surface area and aspect ratio,

respectively. Also, the tendency of the fruit shape towards a sphere can be well determined by the sphericity (Mohsenin, 1978).

Average of 100 fruit mass for the studied variety was 306.54 g. It was estimated 3.06g as average of the individual fruit mass. Turkoglu et al. (2005) reported the hawthorn genotypes fruit mass and flesh/seed ratio as between 4.82 and 9.69 g and 0.29 and 4.21. Higher fruit mass with higher flesh ratio is the most important desirable characteristics of the fruit for breeding programs (Ercisli, 2004).

True density and bulk density of the fruit were 0.89 and 0.47 kg/m<sup>3</sup>, respectively, and porosity was 47.50 %.

A summary of the studied nutritional properties of hawthorn fruit was shown in Table 2.

**Table 2 Nutritional properties of hawthorn fruit**

Parameter	Maximum	Mean	Minimum	Standard Deviation
Moisture content, %	73.10	71.83	69.80	1.78
Dry matter, %	26.10	25.73	25.40	0.35
Total soluble solid (TSS, °Brix)	19.00	18.70	18.3	0.36
Titrateable acidity, %	1.87	1.71	1.62	0.14
TSS/TA ratio	11.60	10.97	9.78	1.03

The moisture content of the hawthorn was 71.83%, w.b.; consequently its dry matter was 25.73%. The moisture content is very important when determining the physical properties such as bulk density, fruit density, porosity, pulp mass, static and dynamic coefficient of friction of fruits and vegetables (Ozcan et al., 2005).

The Total soluble solid (TSS, °Brix) value derived from hawthorn juice was 18.70. Eventually, titratable acidity of the hawthorn fruit was calculated as 1.71%. Using these data, the TSS/TA ratio of the fruit was 10.97.

In a similar research, Ozcan et al. (2005) reported 14.39 mm as fruit length, 19.34 mm as fruit diameter, 3.03 g as fruit mass, 0.87 g as seed mass, 1.98% as acidity and 32.31% as soluble solids content for fresh wild hawthorn fruits collected from Turkey. Turkoglu et al. (2005) recorded fruit diameter between 15.15 and 18.60 mm and fruit mass between 2.24 and 2.44 g for different genotypes of *C. orientalis*. Sparks and Martin (1999) determined fruit mass of *C. monogyna* from 0.21 g to 0.34 g. In next work, Balta et al. (2006) found that among *Crataegus* species, *C. tanacetifolia* had the highest fruit mass (4.99 g) followed by *C. orientalis* (3.48 g), *C. pontica* (3.31 g), *C. aronia* (2.63 g) and *C. meyeri* (1.36 g), respectively. In other researches, conducted by Balta et al. (2006), Ozcan et al. (2005), and Turkoglu et al. (2005) in different regions of Turkey, the total soluble solids and pH values of hawthorn genotypes ranged from 11.66% to 24.00% and 3.12% to 4.09%, respectively.

The studied properties of hawthorn in the presented research can be compared with apple fruits. As Kheiralipour (2008) and Ragni and Berardinelli (2001) reported, the TSS value of hawthorn is much higher than apple fruit. The TSSs of apple fruits were 12.54, 10.73, 14.3, 13.7, 14.3 and 12.9 °Brix for TSS for Delbarstival and Redspar, Golden Delicious, Stark Delicious, Grani Smith and Rome Beauty variety, respectively.

Also value of titratable acidity for the hawthorn is higher than that of apple. The reported data for TSS/TA

ratio of apples were so higher values as 429.20 and 305.85 for Redspar and Delbarstival, respectively (Kheiralipour, 2008).

## 6 Conclusions

Hawthorn is a forest tree grown in different temperate regions of the world. Fresh fruits need to be harvested, transported, sorted, graded and packed. As a first step in design of specific equipment for these processes, physical properties of the fruit need to be known. Some engineering properties of hawthorn fruit harvested from Ilam forests including physical characteristics and nutritional properties were determined to introduce the fruit. Also, the corresponding data those are important for the design of equipments for harvesting, processing, transportation, sorting, separation, packaging and further breeding efforts of hawthorn.

## References

- Bahorun, T., E. Aumjaud, H. Ramphul, M. Rycha, A. Luximon-Ramma, F. Trotin, and O. I. Aruoma. 2003. Phenolic constituents and antioxidant capacities of *Crataegus monogyna* (Hawthorn) callus extracts. *Nahrung/Food*, 47(3): 191–198.
- Balta, M. F., F. Celik, N. Turkoglu, K. Ozrenk, and F. Ozgokce. 2006. Some fruit traits of hawthorn (*crataegus* spp.) genetic resources from malatya, Turkey. *Research Journal of Agriculture and Biological Sciences*, 2(6): 531-536.
- Blumenthal, M., A. Goldberg, and J. Brinckmann. 2000. Hawthorn in herbal medicine expanded: commission e monographs. American Botanical Council, Integrative Medicine Communications, Newton, MA, 182–189.
- Donmez, A. A. 2004. The genus *Crataegus* L. (Rosaceae) with special reference to hybridisation and biodiversity in Turkey, *Turkish J Bot*, 28(1-2): 29–37.
- Ercisli, S. 2004. A short review of the fruit germplasm resources of Turkey. *Genet Resour Crop Evol*, 51(4): 419–435.
- Fadavi, A., S. R. Hassan-Beygi, F. and Karimi. 2013. Moisture dependent physical and mechanical properties of Syrjan region wild pistachio nut. *Agricultural Engineering International: CIGR Journal*, 15(2): 221-230.
- Ghasemi Varnamkhasti, M., H. Mobli, A. Jafari, A. R. Keyhani, M. H. Soltanabadi, S. Rafiee, and K. Kheiralipour. 2008. Some physical properties of rough rice (*Oryza Sativa* L.) grain. *Journal of Cereal Science*, 47(3): 496-501.

- Güven, K., E. Yücel, and F. Çetintas. 2006. Antimicrobial activities of fruits of *Crataegus* and *Pyrus* species. *Pharmaceutical Biology*, 44(2): 79-83.
- Kabas, O., E. Yılmaz, A. Özmerzi, and İ. Akinci. 2007. Some physical and nutritional properties of cowpea seed (*Vigna sinensis* L.). *Journal of food engineering*, 79(4): 1405-1409.
- Kheiralipour, K. 2008. Determination of terminal velocity of two varieties apple (Redspar and Delbarstival) using water column. Un published M.Sc. thesis, University of Tehran.
- Koocheki, A., S. M. A. Razavi, E. Milani, T. M. Moghadam, M. Abedini, S. Alamatian, and S. Izadkhah. 2007. Physical properties of watermelon seed as a function of moisture content and variety. *International Agrophysics*, 21(4): 349.
- Kumar, D., V. Arya, Z. Ali Bhat, N. A. Khan, and D. N. Prasad. 2012. The genus *Crataegus*: chemical and pharmacological perspectives. *Brazilian Journal of Pharmacognosy*, 22(5): 1187-1200.
- Milani, E., M. Seyed, A. Razavi, A. Koocheki, V. Nikzadeh, N. Vahedi, M., Moein Fard, and A. GholamhosseinPour. 2007. Moisture dependent physical properties of cucurbit seeds. *International Agrophysics*, 21(2): 157-168.
- Mirzabe, A. H., J. Khazaei, G. R. Chegini, and O. Gholami. 2013. Some physical properties of almond nut and kernel and modeling dimensional properties. *Agricultural Engineering International: CIGR Journal*, 15(2): 256-265.
- Mohsenin, N. N. 1986. Physical properties of plant and animal materials. Gordon and Breach Press, New York.
- Omobuwajo, O. T., Akande, A. E and Sanni A. L. 1999. Selected physical, mechanical and aerodynamic properties African breadfruit (*Treculia Africana*) seeds. *Food. Eng.*, 40(4): 241-244.
- Ozcan, M., H. Haciseferogullari, T. Marakoglu, and D. Arslan. 2005. Hawthorn (*Crataegus* spp.) fruit: some physical and chemical properties. *J Food Engineering*, 6(4): 409-413.
- Ozturk, N., and M. Tuncel. 2011. Assessment of phenolic acid content and In vitro antiradical characteristics of hawthorn. *Journal of Medical Food*, 14(6): 664-669.
- Perez, E. E., G. H. Crapiste, and A. A. Carelli. 2007. Some physical and morphological properties of wild sunflower seeds. *Biosystems engineering*, 96(1): 41-45.
- Ragni, L., and A. Berardinelli. 2001. Mechanical behavior of apples and damage during sorting and packaging. *J. Agric. Res.*, 78(3): 273-279.
- Riyahi, R., S. Rafiee, M. J. Dalvand, and A. Keyhani. 2011. Some physical characteristics of pomegranate, seeds and arils. *Journal of Agricultural Technology*, 7(6): 1523-1537.
- Sparks, T. H. and T. Martin. 1999. Yields of hawthorn *Crataegus monogyna* berries under different hedgerow management. *Agriculture, Ecosystems and Environment*, 72(2):107-110.
- Turkoglu, N., Kazankaya, and A. Sensoy, R. I. 2005. Pomological characteristics of hawthorn species found in Van Region. *J Agric. Sci*, 15(1): 17-21.
- Veveris, M., Koch, E. and Chatterjee, S. S. 2004. *Crataegus* special extract WS1442 improves cardiac function and reduces infarct size in a rat model of prolonged coronary ischemia and reperfusion. *Life Sci*, 74(15):1945-1955.